

Nakamura T, Okada T, Endo M, Nakamura T, Oike Y, Mizuta H:

Angiopoietin-like protein 2 promotes inflammatory conditions in the ligamentum flavum in the pathogenesis of lumbar spinal canal stenosis by activating interleukin-6 expression.

Eur Spine J 24:2001-2009, 2015

Fujiyuki T, Yoneda M, Amagai Y, Obayashi K, Ikeda F, Shoji K, Murakami Y, Sato H, Kai C.

A measles virus selectively blind to signaling lymphocytic activation molecule shows anti-tumor activity against lung cancer cells.

Oncotarget 6:24895-24903, 2015

Tanaka Y, Ishitsuka Y, Hayasaka M, Yamada Y, Miyata K, Endo M, Kondo Y, Moriuchi H, Irikura M, Tanaka KI, Mizushima T, Oike Y, Irie T:

The exacerbating roles of CCAAT/enhancer-binding protein homologous protein (CHOP) in the development of bleomycin-induced pulmonary fibrosis and the preventive effects of tauroursodeoxycholic acid (TUDCA) against pulmonary fibrosis in mice.

Pharmacol Res 99:52-62, 2015

Masuda T, Endo M*, Yamamoto Y, Odagiri H, Kadomatsu T, Nakamura T, Tanoue H, Ito H, Yugami M, Miyata K, Morinaga J, Horiguchi H, Motokawa I, Terada K, Morioka MS, Manabe I, Iwase H, Mizuta H, Oike Y (***corresponding author**):

ANGPTL2 increases bone metastasis of breast cancer cells through enhancing CXCR4 signaling.

Sci Rep 5:9170, 2015

Aoi J, Endo M*, Kadomatsu T, Miyata K, Ogata A, Horiguchi H, Odagiri H, Masuda T, Fukushima S, Jinnin M, Hirakawa S, Sawa T, Akaike T, Ihn H, Oike Y (***corresponding author**):

Angiopoietin-like protein 2 accelerates carcinogenesis by activating chronic inflammation and oxidative stress.

Mol Cancer Res 12:239-249, 2014

Nakamura T, Okada T, Endo M, Kadomatsu T, Taniwaki T, Sei A, Odagiri H, Masuda T,

Fujimoto T, Nakamura T, Oike Y, Mizuta H:

Angiopoietin-like protein 2 induced by mechanical stress accelerates degeneration and hypertrophy of the ligamentum flavum in lumbar spinal canal stenosis.

PLoS One e85542, 2014

Odagiri H, Kadomatsu T, Endo M, Masuda T, Morioka M, Fukuhara S, Miyamoto T, Kobayashi E, Miyata K, Aoi J, Horiguchi H, Nishimura N, Terada K, Yakushiji T, Manabe I, Mochizuki N, Mizuta H, Oike Y:

The Secreted Protein ANGPTL2 Promotes Metastasis of Osteosarcoma Cells Through Integrin $\alpha 5\beta 1$, p38 MAPK, and Matrix Metalloproteinases.

Sci Signal 7:ra7, 2014

Horio E, Kadomatsu T, Miyata K, Arai Y, Hosokawa K, Doi Y, Ninomiya T, Horiguchi H, Endo M, Tabata M, Tazume H, Tian Z, Takahashi O, Terada K, Takeya M, Hao H, Hirose N, Minami T, Suda T, Kiyohara Y, Ogawa H, Kaikita K, Oike Y:

Role of Endothelial Cell-Derived Angptl2 in Vascular Inflammation Leading to Endothelial Dysfunction and Atherosclerosis Progression.

Arterioscler Thromb Vasc Biol 34:790-800, 2014

Endo M*, Yamamoto Y, Nakano M, Masuda T, Odagiri H, Horiguchi H, Miyata K, Kadomatsu T, Motokawa I, Okada S, Iwase H, Oike Y (*corresponding author):

Serum ANGPTL2 levels reflect clinical features of breast cancer patients: implications for the pathogenesis of breast cancer metastasis.

Int J Biol Markers 29:e239-e245, 2014

Swain L, Wottawa M, Hillemann A, Beneke A, Odagiri H, Terada K, Endo M, Oike Y, Farhat K, Katschinski DM:

Prolyl-4-hydroxylase domain 3 (PHD3) is a critical terminator for cell survival of macrophages under stress conditions.

J Leukoc Biol 96:365-375, 2014

Horiguchi H, Endo M*, Miyamoto Y, Sakamoto Y, Odagiri H, Masuda T, Kadomatsu T, Tanoue H, Motokawa I, Terada K, Morioka MS, Manabe I, Baba H, Oike Y

(*corresponding author):

Angiopoietin-like protein 2 renders colorectal cancer cells resistant to chemotherapy by activating spleen tyrosine kinase-phosphoinositide 3-kinase-dependent anti-apoptotic signaling.

Cancer Sci 105:1550-1559, 2014

Kadomatsu T, Uragami S, Akashi M, Tsuchiya Y, Nakajima H, Nakashima Y, Endo M, Miyata K, Terada K, Todo T, Node K, Oike Y:

A molecular clock regulates Angiopoietin-like protein 2 expression.

PLoS ONE 8: e57921, 2013

Ogata A, Endo M*, Aoi J, Takahashi O, Kadomatsu T, Miyata K, Tian Z, Jinnin M, Fukushima S, Ihn H, Oike Y (*corresponding author):

The role of Angiopoietin-like protein 2 in pathogenesis of dermatomyositis.

Biochem Biophys Res Commun 418:494-499, 2012

Endo M, Nakano M, Kadomatsu T, Fukuhara S, Kuroda H, Mikami S, Hato T, Aoi J, Horiguchi H, Miyata K, Odagiri H, Masuda T, Harada M, Horio H, Hishima T, Nomori H, Ito T, Yamamoto Y, Minami T, Okada S, Takahashi T, Mochizuki N, Iwase H, Oike Y:

Tumor cell-derived angiopoietin-like protein ANGPTL2 is a critical driver of metastasis.

Cancer Res 72: 1784-1794. 【Selected for cover page, April, 2012】 , 2012

Tazume H, Miyata K, Tian Z, Endo M, Horiguchi H, Takahashi O, Horio E, Tsukano H, Kadomatsu T, Nakashima Y, Kunitomo R, Kaneko Y, Moriyama S, Sakaguchi H, Okamoto K, Hara M, Yoshinaga T, Yoshimura K, Aoki H, Araki K, Hao H, Kawasuji M, Oike Y:

Macrophage-derived Angiopoietin-like Protein 2 Accelerates Development of Abdominal Aortic Aneurysm.

Arterioscler Thromb Vasc Biol 32:1400-1409, 2012

Doi T, Kanai T, Mikami Y, Sujino T, Jun L, Ono Y, Hayashi A, Hibi T:

IgA plasma cells express the negative regulatory co-stimulatory molecule programmed cell death 1 ligand and have a potential tolerogenic role in the intestine.

Biochem Biophys Res Commun 425:918-923, 2012

Mizukami T, Kanai T, Mikami Y, Hayashi A, Doi T, Handa T, Matsumoto A, Jun L, Matsuoka K, Sato T, Hisamatsu T, Hibi T:

CCR9+ macrophages are required for eradication of peritoneal bacterial infections and prevention of polymicrobial sepsis.

Immunol Lett 2012 147:75-79, 2012

Kato L, Begum NA, Burroughs AM, Doi T, Kawai J, Daub CO, Kawaguchi T, Matsuda F, Hayashizaki Y, Honjo T:

Nonimmunoglobulin target loci of activation-induced cytidine deaminase (AID) share unique features with immunoglobulin genes.

Proc Natl Acad Sci USA 109:2479-2484, 2012

Miyazaki Y, Kaikita K, Endo M, Horio E, Miura M, Tsujita K, Hokimoto S, Yamamuro M, Iwawaki T, Gotoh T, Ogawa H, Oike Y:

C/EBP Homologous Protein Deficiency Attenuates Myocardial Reperfusion Injury by Inhibiting Myocardial Apoptosis and Inflammation.

Arterioscler Thromb Vasc Biol 31:1124-1132, 2011

Muramoto A, Tsushita K, Kato A, Ozaki N, Tabata M, Endo M, Oike Y, Oiso Y:

Angiotensin-like protein 2 sensitively responds to weight reduction induced by lifestyle intervention on overweight Japanese men.

Nutrition & Diabetes 1: e20, 2011

Aoi J, Endo M*, Kadomatsu T, Miyata K, Nakano M, Horiguchi H, Ogata A, Odagiri H, Yano M, Araki K, Jinnin M, Ito T, Hirakawa S, Ihn H, Oike Y (***First two authors are equal contribution**):

Angiotensin-like Protein 2 is an Important Facilitator of Inflammatory Carcinogenesis and Metastasis.

Cancer Res 71: 7502-7512 [Selected for cover page, December, 2011] , 2011

Sujino T, Kanai T, Ono Y, Mikami Y, Hayashi A, Doi T, Matsuoka K, Hisamatsu

T, Takaishi H, Ogata H, Yoshimura A, Littman DR, Hibi T.

Regulatory T cells suppress development of colitis, blocking differentiation of T-helper 17 into alternative T-helper 1 cells.

Gastroenterology 141:1014-1023, 2011

Nakayama Y, Endo M*, Tsukano H, Mori M, Oike Y, Gotoh T (***First two authors are equal contribution**):

Molecular Mechanisms of the LPS-induced Non-apoptotic ER Stress-CHOP Pathway.

J Biochem 147:471-483, 2010

Okada T, Tsukano H, Endo M, Tabata M, Miyata K, Kadomatsu T, Miyashita K, Semba K, Nakamura E, Tsukano M, Mizuta H, Oike Y:

Synoviocyte-derived angiopoietin-like protein 2 contributes to synovial chronic inflammation in rheumatoid arthritis.

Am J Pathol 176:2309-2319, 2010

Tsukano H, Gotoh T, Endo M, Miyata K, Tazume H, Kadomatsu T, Yano M, Iwawaki T, Kohno K, Araki K, Mizuta H, Oike Y:

The endoplasmic reticulum stress-C/EBP homologous protein pathway-mediated apoptosis in macrophages contributes to the instability of atherosclerotic plaques.

Arterioscler Thromb Vasc Biol 30:1925-1932, 2010

Hato T, Kimura Y, Morisada T, Koh GY, Miyata K, Tabata M, Kadomatsu T, Endo M, Urano T, Arai F, Araki K, Suda T, Kobayashi K, Oike Y:

Angiopoietins contribute to lung development by regulating pulmonary vascular network formation.

Biochem Biophys Res Commun 381:218-223, 2009

Namba T, Tanaka K, Ito Y, Ishihara T, Hoshino T, Gotoh T, Endo M, Sato K, Mizushima T: Positive role of CCAAT/enhancer-binding protein homologous protein, a transcription factor involved in the endoplasmic reticulum stress response in the development of colitis.

Am J Pathol 174:1786-1798, 2009

Shinriki S, Jono H, Ota K, Ueda M, Kudo M, Ota T, Oike Y, Endo M, Ibusuki M, Hiraki A, Nakayama H, Yoshitake Y, Shinohara M, Ando Y:

Humanized anti-interleukin-6 receptor antibody suppresses tumor angiogenesis and in vivo growth of human oral squamous cell carcinoma.

Clin Cancer Res 15:5426-5434, 2009

Tabata M, Kadomatsu T, Fukuhara S, Miyata K, Ito Y, Endo M, Urano T, Zhu HJ, Tsukano H, Tazume H, Kaikita K, Miyashita K, Iwawaki T, Shimabukuro M, Sakaguchi K, Ito T, Nakagata N, Yamada T, Katagiri H, Kasuga M, Ando Y, Ogawa H, Mochizuki N, Itoh H, Suda T, Oike Y:

Angiopoietin-like protein 2 promotes chronic adipose tissue inflammation and obesity-related systemic insulin resistance.

Cell Metab 10:178-188, 2009

Kobayashi M, Aida M, Nagaoka H, Begum NA, Kitawaki Y, Nakata M, Stanlie A, Doi T, Kato L, Okazaki IM, Shinkura R, Muramatsu M, Kinoshita K, Honjo T:

AID-induced decrease in topoisomerase 1 induces DNA structural alteration and DNA cleavage for class switch recombination.

Proc Natl Acad Sci USA 106:22375-22380, 2009

Doi T, Kato L, Ito S, Shinkura R, Wei M, Nagaoka H, Wang J, Honjo T:

The C-terminal region of activation-induced cytidine deaminase is responsible for a recombination function other than DNA cleavage in class switch recombination.

Proc Natl Acad Sci USA 106:2758-2763, 2009

Begum NA, Stanlie A, Doi T, Sasaki Y, Jin HW, Kim YS, Nagaoka H, Honjo T:

Further evidence for involvement of a noncanonical function of uracil DNA glycosylase in class switch recombination.

Proc Natl Acad Sci USA 106:2752-2757, 2009

Nonaka T, Doi T, Toyoshima T, Muramatsu M, Honjo T, Kinoshita K:

Carboxy-terminal domain of AID required for its mRNA complex formation in vivo.

Proc Natl Acad Sci USA 106:2747-2751, 2009

Shivarov V, Shinkura R, Doi T, Begum NA, Nagaoka H, Okazaki IM, Ito S, Nonaka T, Kinoshita K, Honjo T:

Molecular mechanism for generation of antibody memory.

Philos Trans R Soc Lond B Biol Sci 364:569-575, 2009

Doi T, Obayashi K, Kadowaki T, Fujii H, Koyasu S. PI3K is a negative regulator of IgE production.

Int Immunol 20:499-508, 2008

Suyama K, Ohmuraya M, Hirota M, Ozaki N, Ida S, Endo M, Araki K, Gotoh T, Baba H, Yamamura K:

C/EBP homologous protein is crucial for the acceleration of experimental pancreatitis.

Biochem Biophys Res Commun 367:176-182, 2008

Shiroki F, Matsuda S, Doi T, Fujiwara M, Mochizuki Y, Kadowaki T, Suzuki H, Koyasu S: The p85alpha regulatory subunit of class IA phosphoinositide 3-kinase regulates beta-selection in thymocyte development.

J Immunol 178:1349-1356, 2007

Obayashi K, Doi T, Koyasu S:

Dendritic cells suppress IgE production in B cells.

Int Immunol 19:217-226, 2007

Endo M, Mori M, Akira S, Gotoh T:

C/EBP homologous protein (CHOP) is crucial for the induction of caspase-11 and the pathogenesis of lipopolysaccharide-induced inflammation.

J Immunol 176:6245-6253, 2006

Endo M, Oyadomari S, Suga M, Mori M, Gotoh T:

The ER stress pathway involving CHOP is activated in the lungs of LPS-treated mice.

J Biochem 138:501-507 [Selected for cover page, October, 2005] , 2005

Endo M, Oyadomari S, Terasaki Y, Takeya M, Suga M, Mori M, Gotoh T:
Induction of arginase I and II in bleomycin-induced fibrosis of mouse lung.
Am J Physiol Lung Cell Mol Physiol 285:L313-L321, 2003

Doi T, Kinoshita K, Ikegawa M, Muramatsu M, Honjo T:
De novo protein synthesis is required for the activation-induced cytidine deaminase
function in class-switch recombination.
Proc Natl Acad Sci USA 100:2634-2638, 2003